

THE AMERICAN JOURNAL OF

MEDICINE & SPORTS

FOR PRIMARY CARE PHYSICIANS

*Peer Reviewed Clinical Papers on
Medical Problems Affecting Participation in Sports*

VOLUME IV NUMBER II

MARCH/APRIL 2002

EDITORIAL

Triggers of Acute Cardiac Events: New Insights

Murray A. Mittleman, MD, DrPH

COMMENTARY

Health Clubs Need Automated External Defibrillators

Barry A. Franklin, PhD

ORIGINAL PAPERS

The Impact of Creatine Supplementation on Anaerobic Performance: A Meta-Analysis

Mark Misic, EdD; George A. Kelley, DA

Women's Actions Related to Health Behaviors After Receiving Bone Mineral Density Results: An Exploratory Study

Sharon M. Nickols-Richardson, PhD, RD
Courtney E. Quinn, MS; Eric K. Clymer, DO

Implementation of a Novel Cyclic Exercise Protocol in Healthy Women

Rochelle L. Goldsmith, EdD; Irving Dardik, MD
Daniel M. Bloomfield, MD; Sean Hagberg, PhD
Stanley Reisman, PhD; Herbert Benson, MD
Joseph E. Mietus, BS; Ian Harnik, BS
Ary L. Goldberger, MD

REVIEW

Increasing Physical Activity in the Obstructive Sleep Apnea Patient

Anthony S. Kaleth, MS; Thomas W. Chittenden, MA
Jennifer S. Blewins, PhD; Brian J. Hawkins, MS
John M. Gregg, DDS; Don Zedalis, MD
William G. Herbert, PhD

COMMENTARIES

Work-Up of the Athlete With Concussion

Robert C. Cantu, MD

Exertional Rhabdomyolysis: Myths and Madness

Priscilla M. Clarkson, PhD

The Physiologic Basis for the Warm-Up in Therapeutic Exercise Programs

Carl Foster, PhD; John P. Porcari, PhD

DEPARTMENTS

Perceptions

Bengt Saltin, MD
John M. Weiler, MD; William W. Storms, MD

Ask the Specialist

David Wang, MD, MS

Exercise as Medicine

David P. Swain, PhD; Kyle J. McInnis, ScD

Abstracts From the World Literature

Ross E. Andersen, PhD

Senior Section

Jonathan R. Mallen, MD; Joshua A. Strassberg, MD
Steven B. Zelicof, MD, PhD

New Therapy Update Bextra

Constance M. Lebrun, MD

Book Reviews

Henry S. Miller, MD

NEW FEATURE

Outreach News From
the American College
of Sports Medicine

An Official Journal of

 **PTP** PROFESSIONAL
TEAM PHYSICIANS
The Champions of Sports Medicine

Implementation of a Novel Cyclic Exercise Protocol in Healthy Women

Rochelle L. Goldsmith, EdD;¹ Irving Dardik, MD;² Daniel M. Bloomfield, MD;³ Sean Hagberg, PhD;² Stanley Reisman, PhD;⁴ Herbert Benson, MD;⁵ Joseph E. Mietus, BS;⁶ Ian Harnik, BS;¹ Ary L. Goldberger, MD⁶

Division of Circulatory Physiology, Columbia University, College of Physicians and Surgeons, New York, NY;¹ Lifewaves International, Califon, NJ;² Division of Cardiology, Columbia University, College of Physicians and Surgeons, New York, NY;³ The Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ;⁴ The Mind/Body Medical Institute, CareGroup, Department of Medicine, Beth Israel Deaconess Medical Center/Harvard Medical School, Boston, MA;⁵ and The Margret and H.A. Rey Institute for Nonlinear Dynamics in Physiology and Medicine and Beth Israel Deaconess Medical Center/Harvard Medical School, Boston, MA⁶

Traditional exercise regimens are based on maintaining a prolonged increase in heart rate, followed by a single recovery period. The authors tested the efficacy of a cyclic exercise protocol designed to generate a series of parabolic-like waves of cardioacceleration (lasting ≤ 1 minute) followed by recovery to a steady state. In an observational study, they studied the effects of this type of cyclic regimen, consisting of four to seven cycles per session, in a group of healthy women ($n=10$; 32–58 years). Cardiorespiratory fitness, autonomic function, and quality of life were assessed before and after 8 weeks of exercise, performed three days per week. Observations were an increase in peak Vo_2 ($p<0.001$) and the ventilatory breakpoint ($p<0.001$), a decrease in resting diastolic blood pressure ($p<0.05$), an increase in heart rate variability during paced breathing ($p<0.05$), as well as a trend toward increased general positive affect ($p<0.06$). The authors conclude that even very short (8-week) implementation of a cyclic exercise protocol involving ≤ 66 minutes of exercise per month may have beneficial effects. Further studies are indicated to compare cyclic and traditional exercise protocols, both in healthy subjects and in selected patient groups. (*Am J Med Sports*. 2002;4: 135–141, 151) ©2002 Le Jacq Communications, Inc.

Address for correspondence:

Rochelle Goldsmith, EdD, Division of Circulatory Physiology, Columbia University College of Physicians and Surgeons, 630 West 168th Street, New York, NY 10032
E-mail: rlg11@columbia.edu
Manuscript received December 17, 2001; accepted January 21, 2002

Traditional aerobic exercise regimens are based on increasing heart rate (HR) to a target range determined by age and fitness and maintaining that level of activity for a sustained period, followed by a “cool-down” phase. Another approach to training, particularly among athletes, employs interval-type protocols in which there is typically a preset, alternating sequence of exercise and rest periods.

In contrast to these conventional routines, we investigated the implementation of a “cyclic” exercise protocol designed to train not only the activation (arousal) phase, but also the relaxation (recovery) phase of exercise in a wave-like or pulsatile fashion. The physiologic importance of “training recovery” is derived from the concept that recovery itself is a dynamic process, mediated primarily by the parasympathetic nervous system.¹ Therefore, training recovery as well as activation may result in greater neuroautonomic plasticity, which, in turn, may enhance healthy function.¹

The cycle protocol described here is tailored to measurements of an individual’s HR during very short (≤ 1 minute) periods of exercise, followed by variable periods of relaxation to a steady state. This preliminary observational study assessed the effects of a short (8-week) course of cyclic exercise

on selected measures of cardiovascular fitness and subjective perception of psychological well-being in healthy adult women.

Methods

SUBJECTS. Subjects were recruited from the nursing staff of Hunterdon Medical Center (Flemington, NJ). Candidates aged 20–70 years were eligible unless their history revealed any of the following: smoking, chronic disease, chronic drug or medication use, musculoskeletal limitations to exercise, or participation in a structured exercise program. Each subject signed informed consent forms approved by the Columbia University and Hunterdon Institutional Review Boards prior to enrollment.

STUDY DESIGN. Participants underwent a baseline evaluation within 1 week prior to an 8-week cyclic exercise regimen and then again within 1 week after completing the program. The evaluation consisted of a maximal oxygen consumption (V_{O_2}) test, measurement of HR variability (HRV) during controlled respiration, and a standard questionnaire to assess subjective sense of well-being. Prior to beginning the exercise regimen, a baseline cycles test (described below) was used to determine each subject's exercise prescription. The cycles test was repeated after the 8-week training period.

CYCLIC EXERCISE PROTOCOL. Overview. The cyclic exercise protocol was designed to generate a series of parabolic-like waves of cardiovascular exercise and recovery.² A single cycle consisted of a short (≤ 1 minute) burst of exercise followed by a period of cardiovascular recovery. Subjects were taught to use a specific relaxation response³ to standardize and facilitate the recovery period. This type of meditative response has been shown to acutely augment parasympathetic nervous system activity⁴ and thus may potentiate the increase in parasympathetic nervous system activity during recovery. Cycles were performed consecutively in sets of four to seven and at specific times of the day. The study was designed

such that cycles to be performed in the afternoon required more exertion than those in morning sessions. This design was motivated by the findings that work performance typically increases in the afternoon.^{5–7} The exercise phase of the cycles consisted of jogging in place on a trampoline. Two of the subjects complained of knee pain during the trampoline exercise. After the first week of training, one of these subjects instead performed her cycles training on an exercise bicycle (Schwinn Airdyne, Boulder, CO), while the other alternated between the trampoline and bicycle. Five-second averaged HRs were monitored and recorded continuously using a Polar NV HR monitor watch and chest strap (Polar Electro Inc., Woodbury, NY) and then downloaded to a computer. Subjects began a prescribed course of cyclic exercise three times weekly for 8 weeks, monitored by a trainer. After the 8 weeks of cycles training, the subjects performed another cycles exercise test.

Cycles Testing. Subjects were requested to refrain from caffeine and meals for 3 hours prior to testing. Upon arrival, subjects were briefed on the five-cycle baseline exercise test protocol, familiarized with the equipment, and instructed in the relaxation response.³ During this first cycle of the testing phase, subjects were asked to exercise at an easy pace (3 on the 10-point Borg scale⁸) until the HR stabilized (defined as varying \pm three beats over 15 seconds), which always occurred in less than 1 minute of exertion. The subject then stopped the exercise, took a deep breath, sat down, and began initiating the recovery phase of the cycle, including the relaxation response. When the subject had recovered, again defined as a stable HR (\pm three beats over 15 seconds), the subject again took a deep breath and held it for 5 seconds. The deep breath further enhances the HR recovery (Figure 1) via a vagally mediated effect.⁹ The subjects performed cycles two, three, and four in a similar manner but increased the level of exertion with each successive cycle. Cycle five, which was used to determine the peak HR for the subsequent sessions, consisted of a very vigorous, brief exertion (9 on the 10-point Borg Scale), continued until the HR

Table I. Cyclic Exercise Protocol: Overview

WEEK	TIME OF DAY	HR RANGES (PEAK HR)	CYCLE SET DAY 1 MONDAY	CYCLE SET DAY 2 WEDNESDAY	CYCLE SET DAY 3 FRIDAY
1	6 a.m.–9 a.m.	60%–92%	7 Cycles	6 Cycles	5 Cycles
2	6 a.m.–9 a.m.	65%–93%	7 Cycles	6 Cycles	5 Cycles
3	9 a.m.–noon	70%–97%	5 Cycles	5 Cycles	4 Cycles
4	3 p.m.–6 p.m.	75%–100%	6 Cycles	5 Cycles	5 Cycles

Peak heart rate (HR) as determined from the baseline cycles exercise test

plateaued or the subject completed 1 minute of exertion. The subjects then began the final recovery cycle. The trainer measured the subject's blood pressure 5 minutes after the peak HR was reached and again 10 minutes later ("recovery blood pressure"). A sample of a five-cycle baseline test is shown in Figure 1.

Exercise Training. Once the baseline cycles testing was completed, the subjects began a regimen consisting of four to seven consecutive cycles per day, three days per week for 8 weeks (Table I). A trainer tracked the participants as they moved through the same type of exercise-recovery cycles described above. Target HRs for each cycle were calculated as percentages of the peak HR determined from the baseline session described above. The target HRs for each cycle began at the low end of the range and progressively increased in a step-wise, incremental fashion to the high end of the range (Table I). An example of the actual exercise prescription for one of the subjects (whose baseline exercise test is shown in Figure 1) is shown in Table II. During cycles that required high HRs, all subjects exercising on the trampoline used hand weights (2–3 pounds) to assist in reaching HR targets within the allotted 60 seconds. Each subject's individual exercise protocol was assessed after 4 weeks of exercise. If a subject had exceeded her target HR on the final cycle of the final session by more than five beats per minute, she was moved into the next higher range of exercise HRs for the second month. If not, she repeated the same exercise protocol for the second month.

MAXIMAL CARDIOPULMONARY EXERCISE TEST. Peak VO_2 was measured by an incremental exercise test on a treadmill. Expired gas analysis of each breath was performed continuously during the test with a SensorMedics Vmax 29 (Yorba Linda, CA) metabolic cart. During the test, the work rate increased as a ramp or step function; treadmill protocol choice was made based on each subject's fitness level as determined by a pretest questionnaire. Peak VO_2 was defined as the highest VO_2 averaged over 20 seconds achieved during the final minute of exercise. VO_2 at the ventilatory breakpoint (VB) was identified using the V-slope method.¹⁰ VB could be determined in nine of the 10 subjects. Subjects' HRs were compared at 3 minutes and 6 minutes into both the pre- and post-training exercise tests to study the effect of exercise training on submaximal HR. Each subject performed the same protocol on her pre- and post-test.

HR VARIABILITY. Studies were performed in a quiet room. The electrocardiogram (EKG) was monitored on an oscilloscope and continuously recorded using a digital acquisition analysis program (Gould Ponemah version 1.21, Valley View, OH). The subjects were allowed to acclimate to the environment while breathing spontaneously. Subjects were then asked to breathe for




Figure 1. A representative set of five cycles performed during the baseline cycles testing. Peak heart rates achieved were 117, 141, 146, 160, and 176 beats per minute (bpm) for cycles 1 through 5, respectively. After each cycle, the subject recovered by sitting quietly and performing the relaxation response. After the heart rate achieved a steady state, the subject took a 5-second deep breath and then began the next cycle. As seen on the graph (arrow), deep breathing was associated with a further lowering of heart rate during recovery.

7 minutes at a rate of 12 breaths per minute with the aid of a light box. A 6.5-minute window at the start of paced breathing was selected for analysis, this being the longest data length common to all subjects. The mean and standard deviation of the R-R intervals (SDNN) over this window were calculated. Spectral powers were calculated for total power (0–0.40 Hz), low-frequency power (0.04–0.15 Hz), and high-frequency power (0.15–0.40 Hz), using the Lomb periodogram technique for unevenly sampled data.¹¹

PSYCHOLOGICAL MEASUREMENTS. The 38-item Mental Health Inventory (MHI)¹² was used to measure changes in psychological status. The MHI includes six subscales—Anxiety, Depression, Loss of Behavioral/Emotional Control, General Positive Affect, Emotional Ties, and Life Satisfaction; two global scales—Psychologic Distress and Psychologic Well-Being; and an overall Mental Health Index. For each item, subjects rated the frequency or intensity of a psychological symptom during the previous month. All items except two used a six-point response scale; two of the items used five-point response scales.

STATISTICAL ANALYSES. Comparisons of HR, HRV, and fitness measures at baseline and after 8 weeks of

HR cycle training were analyzed using the Student *t* test for paired observations. In order to use parametric statistics, which require near-normal distributions, HRV data were transformed to their natural logarithms. Systolic and diastolic blood pressure (SBP and DBP) (recorded at the time of the cycles exercise test) at rest and during recovery, before and after cycles training, were compared using a two-way repeated

measures analysis of variance. The Wilcoxon signed rank test was used to determine pre- and postprogram differences on all scales of the MHI. Statistical significance is defined as $p < 0.05$.

Results

DESCRIPTIVE DATA. Eleven subjects were recruited for this study. One of the subjects developed nausea and emesis in response to cycles testing during the first set of cycles. This subject did not continue with the prescribed exercise program. The baseline characteristics of the 10 subjects who completed the protocol are shown in Table III. The peak VO_2 levels of the subjects are consistent with those of untrained healthy women.

MAXIMAL CARDIOPULMONARY EXERCISE TEST. Results of treadmill testing at baseline and after completion of the training are shown in Table IV. Peak VO_2 increased by 14.8% ($p < 0.0001$). As seen in Figure 2, nine of the 10 subjects improved their peak VO_2 . VO_2 at the ventilatory breakpoint also increased significantly ($p < 0.003$) after training.

CYCLES TESTING. The peak HR achieved during cycles testing averaged 99% of the peak HR achieved on the maximal treadmill test. Training was not associated with a change in resting SBP (125 ± 14 vs. 124 ± 14 mm Hg) or postexercise recovery SBP (123 ± 15 vs. 125 ± 15 mm Hg). DBP was significantly lower after training in both the pre-exercise resting (79.4 ± 8.4 vs. 73.2 ± 7.9 mm Hg) and postexercise recovery states (73.8 ± 14.8 vs. 68.4 ± 10.2 mm Hg; $p < 0.009$). In addition, DBP decreased approximately 5 mm Hg from resting to recovery phases ($p < 0.04$) in both the trained and untrained states.

HR AND HRV. After 8 weeks of exercise, despite no significant change in mean resting HR, a reduction in mean HR at submaximal workloads was observed. At 3 minutes into the postexercise tests, mean HR was 2.4 ± 3.5 beats/min lower than it had been on the pretests ($p < 0.06$); at 6 minutes, mean HR was 4.7 ± 5.5 beats/min lower ($p < 0.03$). There was also a small but significant increase in HRV after training (Table V). Broadband measures of HRV, such as the SD of normal R-R intervals (SDNN) and the natural logarithm total power, increased ($p < 0.05$). In high-frequency power, a measure of vagally mediated R-R variability, increased 9.1% ($p < 0.05$) when measured during paced breathing.

PSYCHOLOGICAL MEASUREMENTS. There was a trend toward a decrease in anxiety and an increase in general positive affect, psychological well-being, and overall mental health index after 8 weeks of cycles training (Table VI).

Table II. Exercise prescription for a subject who achieved a peak heart rate (HR) of 176 beats/min on her baseline cycles test. The numbers listed below are the HR goals for each cycle.

WEEK 1 6 A.M.–9 A.M.			
Monday	Wednesday	Friday	
107	111	115	
125	131	134	
137	149	148	
147	5-Minute break	153	
5-Minute break	145	160	
143	153		
151	158		
156			
WEEK 2 6 A.M.–9 A.M.			
Monday	Wednesday	Friday	
112	116	120	
130	136	139	
142	154	153	
152	5-Minute break	158	
5-Minute break	150	165	
148	158		
156	163		
161			
WEEK 3 9 A.M.–NOON			
Monday	Wednesday	Friday	
123	126	129	
144	147	152	
152	155	160	
161	164	170	
167	169		
WEEK 4 3 P.M.–6 P.M.			
Monday	Wednesday	Friday	
132	135	138	
147	150	154	
155	162	162	
168	167	167	
159	172	175	
169			
The time windows indicate the hours during the day in which exercise sessions were conducted.			

